

Peragreen® 15%

(ANTIMICROBIAL SOLUTION)

Peragreen® 15% is a peroxyacetic acid-based microbiocide developed for use in Fruit and Vegetable Process Water Systems and for Agricultural or Horticultural uses.

ACTIVE INGREDIENT:

Peroxyacetic Acid 15.0%
Hydrogen Peroxide 22.0%

INERT INGREDIENTS:

63.0%
TOTAL 100.0%

EPA Registration No: 63838-2

EPA Est. No. 63838-CA-01: 63838-AR-001

Before Using This Product, Please Read This Entire Label Carefully.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
QUESTIONS? 1-209-581-9576	Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
NOTE TO PHYSICIAN:	Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER CORROSIVE: Do not enter an enclosed area without proper respiratory protection, or when uncoupling of product transfer hoses. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles, face shield, rubber gloves and protective clothing with long sleeves when handling. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

PHYSICAL OR CHEMICAL HAZARDS:

STRONG OXIDIZING AGENT. CORROSIVE: Mix only with potable water below 140° F. Product must be diluted in accordance with label directions prior to use. This product is not combustible; however, at temperatures exceeding 156°F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to birds, fish and aquatic invertebrates. Caution must be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of the National Pollution Discharge System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage plant authority.

STORAGE AND DISPOSAL

Storage: Never return this product to the original container after it has been removed. Avoid all contaminants, especially dirt, caustic, reducing agents, and metals. Contamination and impurities will reduce shelf life and can induce decomposition. In case of a decomposition, isolate container, spray container with

cool water and dilute this product with large volumes of water. Avoid damage to containers. Keep container closed at all times when not in use. Keep container out of direct sunlight. To maintain product quality, store at temperatures below 86°F.

Procedure for Leak or Spill: Stop leak if this can be done without risk. Shut off ignition sources: no flames, smoking, flares, or spark producing tools. Keep combustible and organic materials away. Flush spilled material with large quantities of water. Undiluted material must not enter confined spaces.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance. If material has been spilled, an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all local, state and Federal environmental laws, rules, regulations, standards, and other requirements. Because acceptable methods of disposal may vary by location, regulatory agencies must be contacted prior to disposal. This product which is to be discarded, must be disposed of as hazardous waste after contacting the appropriate local State or Federal agency to determine proper procedures.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Offer for recycling, if available. Closure as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS

This product can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables for the control of bacteria and fungi in commercial operations and packinghouses.

Batch, Continuous or Spray System Processes: Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method. Add this product to no more than 533 ppm (w/wt) total product (80 ppm residual peroxyacetic acid) to the use solution. This can be accomplished by initially adding 53.3 grams (47.3 mls) of this product per 100 liters of water, or 1.0 fl. oz. per 16.4 gallons of water. The fruits and vegetables can be continuously sprayed (using coarse spray) or submerged (dipped) in the resulting solution. Periodic or continuous additions of this product to maintain the required concentration may be added as necessary. It is also recommended to apply this product during the washing, chilling, or physical cleaning processes, including the roller-spreader, washer or brush washer manifold, dip tank, or sorting processes. Contact time of 60 seconds is recommended to insure efficacy. A potable water rinse is not required.

Fogging: (Not for use in California) For raw agricultural commodities, commercially-applied fogging methods may be used provided the dilution rates of the resultant solution does not exceed those prescribed in this section (1 fl oz per 16.4 gal of water). A potable water rinse, or subsequent antimicrobial wash using this product as prescribed above is necessary after such use. Conventional corrosion-resistant fogging devices are recommended. Vacate the area of all personnel prior to, during and after fogging until the total peroxide concentration is below 1.0 ppm, or there is no strong odor present, characteristic of acetic acid.

AGRICULTURAL or HORTICULTURAL USES

There is a Restricted-Entry-Interval of zero (0) hours after the use of this product. This product must never be mixed or combined with any other pesticide or fertilizer. Upon soil contact this diluted product decomposes rapidly to oxygen, carbon dioxide and water. This product may be harmful to fish if exposed on a continuous basis at concentrations of 1 ppm or more of active peroxyacetic acid. Meter this product into pressurized piping using a plastic or stainless steel injection/backflow device installed far enough upstream from the equipment to ensure thorough mixing. For open flowing bodies of water, apply this product as far upstream as possible to allow adequate mixing prior to the flow entering any larger body of water. If open pouring of this product is required pour product as close to the surface of the water as possible to reduce odor exposure.

Treatment of Agricultural or Irrigation Water Systems (sand filters, humidification systems, storage tanks, ponds, reservoirs, canals): For the control of sulfides, odor, slime and algae in water systems, apply this product at 2-10 ppm active peroxyacetic acid. This feed rate equals 15-75 fl. oz. per 10,000 gallons of water. Repeat dose as necessary to maintain control, which will vary with seasonal conditions. For prevention of algae, some systems may require continuous low level dosing during warm sunny periods (2-5 ppm peroxyacetic acid).

Drip Irrigation Systems: To clean slime and algae from drip system filters, tapes and emitters, meter this product at the rate of 7.5-15 fl. oz. per 1000 gallons of water (10-20 ppm peroxyacetic acid). When required during normal irrigation cycles, use this product at the recommended dose for a minimum of 30 minutes. After an irrigation cycle do not flush the lines.

Greenhouses: This product can be used to suppress/control algae and slime formations in and around greenhouses. For normal use in various process, irrigation or sprinkler water systems, this product may be used at 1:40,000 to 1:5000 dilutions (4-33 ppm as peroxyacetic acid). Heavily fouled systems, such as evaporative coolers or irrigation/drip lines may need shock doses of up to 100 ppm as peroxyacetic acid (1:1,600 dilution).

Soil Applications: Use this product at 16.4-33 fl. oz. per 100 gal of water (220-440 ppm active peroxyacetic acid) for the control of soil-borne diseases such as Fusarium, Phytophthora, Pythium, Verticillium, Thielaviopsis, and Rhizoctonia. This product can be applied by drench, flood, drip or sprinkler irrigation systems. Best results may be obtained by application prior to and during the seeding or transplant operations. Wait one day before inoculating the soil with beneficial microbes.

Foliar Applications: This product may be used to cure or prevent bacterial and fungal diseases on growing agricultural crops. Typical use rates are 5.8-49.5 fl. oz. of this product per 100 gal of water (77-663 ppm active peroxyacetic acid) applied at 30-100 gal of mixed solution per acre of foliage. Curative (or rescue) treatment requires the lower dilution rates, while preventative treatments use the higher dilution rates. Apply curative treatments for 2-3 days and then resume weekly preventative treatments thereafter. Good coverage and wetting of the foliage is required. Not all plant diseases have been tested, but some of the common diseases controlled are: Algae, Alternaria spp., Anthracnose, Aphanomyces, Bacterial Blight, Black Spot, Botrytis (gray mold), Brown Spot, Copper Spot, Dollar Spot, Early and Late Blights, Erwinia spp., Fairy Ring, Fusarium Root Rot and Blight, Fruit, Black, Brown, Stem and Sour Rots, Leaf and Bacterial Spots, Plasmopara, Powdery and Downy mildews, Phytophthora Blight/Rots, Pink Snow Mold, Pseudomonas and Xanthomonas spp., Pythium spp., Rhizoctonia spp., Rusts, Scabs, Scum, Slime Molds, Smut, Summer Patch, Stripe Smut, Take-all Patch, and Thielaviopsis.

Manufactured By:

ENVIRO TECH CHEMICAL SERVICES, Inc.

500 Winmoore Way, Modesto, CA 95358

209-581-9576 or www.envirotech.com

LOT #:

Net contents: 500 lbs. / 52.79 gal.

63-QA100 Ver 9Y

DOT: **UN 3109**, Organic Peroxide Type F, Liquid
(≤25% Peracetic Acid with ≤26% Hydrogen Peroxide) 5.2 (8)